

Temperature & Pressure Conversions WKSHT

- | | |
|-------------------------------------|-----------------------------------------------|
| 1.) 2.00 atm to mm Hg | 9.) 1800. mm Hg to kPa |
| 2.) 115 kPa to atm | 10.) 93,500 Pa to atm |
| 3.) 500. mm Hg to atm | 11.) 950. torr to atm |
| 4.) 3.5×10^4 torr to mm Hg | 12.) 0.490 atm to kPa |
| 5.) 35 °C to Kelvin | 13.) standard temperature in Kelvin & Celsius |
| 6.) 120 °C to Kelvin | 14.) 298 K to °C |
| 7.) -25 °C to Kelvin | 15.) 100. K to °C |
| 8.) -227 °C to Kelvin | 16.) 5 Kelvin to °C |

Ideal Gas Equation 1 WKSHT

- 1.) What is the pressure exerted by 2.0 moles of an ideal gas when it occupies a volume of 12.0 L at 373 K?
- 2.) A flashbulb of volume 2.6 cm^3 contains O_2 gas at a pressure of 2.3 atm and a temperature of 26°C. How many moles of O_2 does the flashbulb contain?
- 3.) If 0.20 moles of helium occupies a volume of 64.0 liters at a pressure of 0.15 atm, what is the temperature of the gas?
- 4.) What is the volume of 0.35 moles of gas at 1.7 atm of pressure and a temperature of 100 K?
- 5.) What is the pressure of 1.5 moles of an ideal gas at a temperature of 150 K and occupies a volume of 20.0 liters?
- 6.) How many moles of gas occupy 16.2 liters at a pressure of 1.05 atm and a temperature of 37°C?

Answers: 1.) 5.1 atm; 2.) 2.4×10^{-4} moles; 3.) 580 K; 4.) 1.7 L; 5.) 0.92 atm; 6.) 0.668 moles

Ideal Gas Equation 2 WKSHT

- 1.) Calculate the volume of exactly 1.00 mole of a gas at STP.
- 2.) How many moles of nitrogen are present in 17.8 liters at 27 °C and 1.3 atm pressure?
- 3.) What is the pressure of 2.3 moles of carbon dioxide at 235 K occupying 23.7 liters of space?
- 4.) If there are 4.02×10^{23} molecules of N_2O in a sample, how many moles are there?
- 5.) Using answer from #6, calculate the pressure of the gas if it occupies $27,025 \text{ cm}^3$ of space at 38.0 °C.
- 6.) How many grams of NH_3 are present in 35.0 dm^3 of space at 78.3 K and 0.853 atm of pressure?
- 7.) What is the temperature of 34.2 grams of sulfur dioxide occupying 30.0 liters of space and having a pressure of 800. torr?
- 8.) What is the pressure (in mm Hg) of 79.4 grams of boron trifluoride in a 20.0 L container at a temperature of 245 K?
- 9.) How many grams are in a sample of arsenic trifluoride that has a volume of 17,600 mL and a temperature of 92 °C and a pressure of 108,732 Pa?
- 10.) How many kilopascals of pressure are exerted by 23.8 liters of oxygen with a mass of 175 grams at a temperature of 58 °C?
- 11.) How many moles of argon are in 30.6 liters at 28 K and 658 mm Hg of pressure?
- 12.) How many grams of argon are found in # 11?

Answers: 1.) 22.4 L; 2.) 0.94 moles; 3.) 1.9 atm; 4.) 0.668 moles; 5.) 0.631 atm; 6.) 78.9 g; 7.) 720. K; 8.) 897 mm Hg; 9.) 83 g; 10.) 632 kPa; 11.) 12 moles; 12.) 480 g

Applications of Ideal Gas Equation WKSHT

- 1.) What pressure is exerted by 1.0 mole of an ideal gas contained in a 1.0 L vessel at 0.0 °C?
- 2.) What is the density of a sample of ammonia gas, NH_3 , if the pressure is 0.928 atm and the temperature is 63.0 °C?
- 3.) Calculate the molar mass of a gas if 4.5 L of the gas at 785 torr and 23.5 °C has a mass of 13.5 grams.
- 4.) 0.453 moles of a gas confined to a 15.0 L container exerts a pressure of 1.24 atm on the walls of the container. What is the temperature of the gas (in °C)?

UNIT 10 - GASES

- 5.) 5.4 grams of carbon dioxide are confined to a 20.0 L container at a temperature of 32.5 °C. What pressure does the gas exert?
- 6.) 2.125 grams of a gas in a 1.25 L container exert a pressure of 0.838 atm at 40.0 °C. What is the molar mass of the gas?
- 7.) To what temperature must 10.0 grams of NH₃ have to be heated in a 15.0 L container in order for it to exert a pressure of 3.50 atm?
- 8.) 2.0×10^{-5} grams of hydrogen gas at 155 °C exert a pressure of 322.5 torr on the walls of a small cylindrical tube. What is the volume of the tube?

Answers: 1.) 22 atm; 2.) 0.572 g/L; 3.) 71 g/mole; 4.) 227 °C; 5.) 0.15 atm; 6.) 52.1 g/mole; 7.) 1090 K; 8.) 8.3×10^{-4} L

Gas Law Problems WKSHT

- 1.) The gas pressure in an aerosol can is 1.5 atm at 25 °C. Assuming that the gas inside obeys the ideal gas equation, what would the pressure be if the can were heated to 450 °C?
- 2.) A pocket of gas is discovered in a deep drilling operation. The gas has a temperature of 480 °C and is at a pressure of 12.8 atm. Assume ideal behavior. What volume of the gas is required to provide 18.0 L at the surface at 1.00 atm and 22 °C?
- 3.) A fixed quantity of gas is compressed at constant temperature from a volume of 368 mL to 108 mL. If the initial pressure was 5.22 atm, what is the final pressure?
- 4.) A gas originally at 15 °C and having a volume of 182 mL is reduced in volume to 82.0 mL while its pressure is held constant. What is its final temperature?
- 5.) At 36 °C and 1.00 atm pressure, a gas occupies a volume of 0.600 L. How many liters will it occupy at 0.0 °C and 0.205 atm?
- 6.) What is the temperature at which 9.87×10^{-2} moles occupies 164 mL at 0.645 atm?
- 7.) Chlorine is widely used to purify municipal water supplies and to treat swimming pool waters. Suppose that the volume of a particular sample of Cl₂ is 6.18 L at 0.90 atm and 33 °C. What volume will the Cl₂ occupy at 107 °C and 0.75 atm?
- 8.) A gas exerts a pressure of 1.5 atm at 27 °C. The temperature is increased to 108 °C with no volume change. What is the gas pressure at the higher temperature?

Answers: 1.) 3.6 atm; 2.) 3.59 L or 3.6 L; 3.) 17.8 atm; 4.) 130 K or 130. K; 5.) 2.59 L or 2.6 L; 6.) 13.1 K; 7.) 9.2 L; 8.) 1.9 atm

GAS STOICHIOMETRY (standard conditions) WKSHT. (assume STP conditions for all reactions)

- 1.) How many liters of oxygen can be formed from the decomposition of 2.00 grams of KClO₃.
$$\underline{\hspace{1cm}} \text{KClO}_3 \rightarrow \underline{\hspace{1cm}} \text{KCl} + \underline{\hspace{1cm}} \text{O}_2$$
- 2.) How many grams of CaCO₃ are required to produce 6.00 L of CO₂?
$$\underline{\hspace{1cm}} \text{CaCO}_3 \rightarrow \underline{\hspace{1cm}} \text{CaO} + \underline{\hspace{1cm}} \text{CO}_2$$
- 3.) Determine the volume of hydrogen gas produced when 0.250 moles of zinc react with excess HCl.
$$\underline{\hspace{1cm}} \text{Zn} + \underline{\hspace{1cm}} \text{HCl} \rightarrow \underline{\hspace{1cm}} \text{ZnCl}_2 + \underline{\hspace{1cm}} \text{H}_2$$
- 4.) How many liters of nitrogen are required to combine with 3.0 L of hydrogen in the following reaction:
$$\underline{\hspace{1cm}} \text{N}_2 + \underline{\hspace{1cm}} \text{H}_2 \rightarrow \underline{\hspace{1cm}} \text{NH}_3$$
- 5.) How many liters of oxygen are needed to combine with 7.0 liters of propane in the following reaction:
$$\underline{\hspace{1cm}} \text{C}_3\text{H}_8 + \underline{\hspace{1cm}} \text{O}_2 \rightarrow \underline{\hspace{1cm}} \text{CO}_2 + \underline{\hspace{1cm}} \text{H}_2\text{O}$$
- 6.) From the following reaction:
$$\underline{\hspace{1cm}} \text{CH}_4 + \underline{\hspace{1cm}} \text{O}_2 \rightarrow \underline{\hspace{1cm}} \text{CO}_2 + \underline{\hspace{1cm}} \text{H}_2\text{O}$$

How many liters of CO₂ are formed from 32.0 grams of CH₄?
- 7.) How many grams of Na are needed to produce 5.0 L of hydrogen?
$$\underline{\hspace{1cm}} \text{Na} + \underline{\hspace{1cm}} \text{H}_2\text{O} \rightarrow \underline{\hspace{1cm}} \text{NaOH} + \underline{\hspace{1cm}} \text{H}_2$$
- 8.) Determine the volume of CO₂ produced from burning 0.750 moles of C.
$$\underline{\hspace{1cm}} \text{C} + \underline{\hspace{1cm}} \text{O}_2 \rightarrow \underline{\hspace{1cm}} \text{CO}_2$$

Answers: 1.) 0.551 L; 2.) 26.8 g; 3.) 5.60 L; 4.) 1.0 L; 5.) 35 L; 6.) 44.8 L; 7.) 10. g; 8.) 16.8 L

DALTON'S LAW & GRAHAM'S LAW WKSHT.

- Determine the partial pressure of each gas in a container with 2.0 moles of N₂, 3.0 moles of O₂, and 7.0 moles of H₂ that has a total pressure of 850 mm Hg. (You will have 3 separate answers for this question.)
- A mixture of nitrogen and oxygen has a total pressure of 730 mm Hg. If the nitrogen has a partial pressure of 420 mm Hg, find the pressure of the oxygen.
- At an altitude of 30,000 ft., the total air pressure is only about 450. mm Hg. If the air is 21.0 % oxygen, what is the partial pressure of oxygen at this altitude?
- A mixture of 3 gases have the following pressures: oxygen = 355 mm Hg, helium = 468 mm Hg, & nitrogen = 560 mm Hg. Find the % of each gas in the mixture.
- Compare the rate of effusion of CH₄ and CO₂. (Give answers to # 5, 6, & 7 to 3 SF's.)
(Your answers for # 5, 6, & 7 should read " ___ effuses ___ times faster than ___.")
- Compare the rate of effusion of helium and nitrogen.
- How much faster does ammonia (NH₃) effuse than HCl?
- An unknown gas effuses 4.0 times faster than O₂. Find the molar mass of the unknown gas.

Answers: 1.) N₂ = 142 mm Hg, O₂ = 213 mm Hg, H₂ = 496 mm Hg 2.) 310 mm Hg 3.) 94.5 mm Hg 4.) O₂ = 25.7%,
He = 33.8 %, N₂ = 40.5 % 5.) CH₄ effuses 1.66 times faster than CO₂. 6.) He effuses 2.65 times faster than N₂.
7.) NH₃ effuses 1.46 times faster than HCl 8.) 2.0 g/mole

UNIT 10 REVIEW WORKSHEET

- Convert the following pressure measurements to atmospheres.
(A) 151.98 kPa (B) 456 mm Hg (C) 912 torr
- What are the conditions for gas measurement at STP?
- The volume of a sample of methane gas measures 350. mL at 27.0 °C and 810. mm Hg. What is the volume (in liters) at -3.0 °C and 650. mm Hg pressure?
- How many grams of nitrogen gas are contained in a 32.6 liter container at 34.4 °C and 579 torr?
- A mixture of four gases in a container exerts a total pressure of 955 mm Hg. In this container, there are 4.50 moles of N₂, 4.25 moles of CO₂, 2.75 moles of H₂, and 2.00 moles of O₂. What is the partial pressure of H₂?
- Compare the rates of effusion of carbon dioxide gas and carbon monoxide gas.
- An unknown gas effuses 1.37 times faster than chlorine gas. What is the molar mass of the unknown gas?
- Given the following unbalanced reaction: ___ C₅H₁₂ + ___ O₂ → ___ CO₂ + ___ H₂O
How many liters of oxygen are needed to produce 45.7 liters of CO₂?
- Given the unbalanced equation: ___ Mg + ___ O₂ → ___ MgO
How many liters of oxygen gas are required to produce 45.8 grams of magnesium oxide?
- An aerosol can contains gases under a pressure of 4.50 atm at 20.0 °C. If the can is left on a hot, sandy beach, the pressure of the gases increases to 4.80 atm. What is the temperature on the beach (in °C)?

ANSWERS:

- 1.) (A) 1.5003 atm (B) 0.600 atm (C) 1.20 atm 2.) 0 °C (or 273 K) & 1 atm 3.) 0.393 L
4.) 27.6 g 5.) H₂ = 195 mm Hg 6.) CO effuses 1.25 times faster than CO₂.
7.) 37.8 g/mole 8.) 73.1 L 9.) 12.8 L 10.) 40.0 °C

UNIT 10 SUMMARY & PRACTICE WORKSHEET

- Convert the following temperatures.
(A) 104 °C to K (B) -3 °C to K (C) 67 K to °C (D) 1671 K to °C
- Convert the following pressures.
(A) 635 torr to atm (B) 104.2 kPa to mm Hg (C) 1.45 atm to Pa
- A gas that effuses 1.19 times slower than nitrogen is added to light bulbs. What is the molecular mass of this unknown gas?
- (A) What is the molecular mass of a 0.2500 g sample of a gas at 99.8°C and 0.9131 atm in a 100.0 cm³ container? (B) What is the gas in the container?
- A small 2.00 L fire extinguisher has an internal pressure of 506.6 kPa at 25°C. What volume of methyl bromide, the fire extinguisher's main ingredient, is needed to fill an empty fire extinguisher at standard pressure if the temperature remains constant?
- If 45.0 g of propane gas burns completely in the following reaction:
$$\text{C}_3\text{H}_{8(g)} + 5 \text{O}_{2(g)} \rightarrow 3 \text{CO}_{2(g)} + 4 \text{H}_2\text{O}_{(g)}$$
then how many liters of carbon dioxide gas will be released if the system is at STP?

UNIT 10 - GASES

7. Air in a closed cylinder is heated from 25°C to 36°C. If the initial pressure is 3.80 atm, what is the final pressure?
8. At what temperature Celsius will 19.4 g of molecular oxygen, O₂, exert a pressure of 1820 mm Hg in a 5.12 L cylinder?
9. To what temperature must 32.0 ft³ of a gas at 2.0 °C be heated for it to occupy 1.00 x 10² ft³ at the same pressure? (ft³ is a unit of volume)
10. Determine the molar mass of a gas that has a density of 2.18 g/L at 66°C and 720 mm Hg.
11. A 3.10 mL bubble of methane gas forms at the bottom of a bog where the temperature is 12°C and the pressure is 8.5 atm. The bubble rises to the surface where the temperature is 35°C and the pressure is 1.18 atm. What is the new volume of the methane bubble?
12. A mixture of 2.00 moles of H₂, 2.00 moles of NH₃, 4.00 moles of CO₂ and 5.00 moles of N₂ exerts a total pressure of 800. torr. What is the partial pressure of the carbon dioxide gas?
13. For the reaction 2 H_{2(g)} + O_{2(g)} → 2 H_{2O(g)}, how many liters of water can be made from 5.0 L of oxygen gas and an excess of hydrogen?

ANSWERS: 1. (A) 377 K (B) 270 K (C) -206 °C (D) 1398 °C
 2. (A) 0.836 atm (B) 781.8 mm Hg (C) 147,000 Pa 3. 39.7 g/mole
 4. (A) 83.8 g/mole (B) Kr 5. 10.0 L 6. 68.5 L 7. 3.94 atm 8. -27 °C
 9. 859 K 10. 64.1 g/mole 11. 24 mL 12. 246 torr 13. 10. L

GASES - A FANTASTIC SUMMARY & REVIEW!

Law	Ideal Gas Equation	Combined Gas Law
equation	$P V = n R T$	$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$
explanation	one gas at one set of conditions	one gas that is changing conditions
when to use it	when the problem gives 3 of these: P, V, n, T	more than one temperature, pressure, and/or volume in the problem
specific units req'd?	pressure = atm, volume = liters, quantity (n) = moles, temperature = Kelvins	temperature = Kelvins pressure & volume can be any unit, but same on both sides of the equation

Law	Boyle's Law	Charles' Law	Gay-Lussac's Law
equation	$P_1 V_1 = P_2 V_2$	$\frac{V_1}{T_1} = \frac{V_2}{T_2}$	$\frac{P_1}{T_1} = \frac{P_2}{T_2}$
explanation	pressure & volume are inversely proportional; temperature is constant	volume & Kelvin temp of a gas are directly proportional; P is constant	pressure & Kelvin temp of a gas are directly proportional; V is constant
when to use it	given 2 different pressures & 1 volume or given 2 different volumes & 1 pressure	given 2 different volumes & 1 temperature or given 2 different temperatures & 1 volume	given 2 different pressures & 1 temperature or given 2 different temperatures & 1 pressure
specific units req'd?	any - but must be the same on both sides of equation	any unit for volume (same on both sides), Kelvin temperature	any unit for pressure (same on both sides), Kelvin temperature

UNIT 10 - GASES

Law	Dalton's Law	Dalton's Law	Graham's Law
equation	$P_{\text{total}} = P_{\text{gas1}} + P_{\text{gas2}} + \dots$	$P_x = \frac{(\text{moles } x)}{(\text{total moles})} \cdot P_{\text{total}}$	$\frac{\text{rate A}}{\text{rate B}} = \sqrt{\frac{\text{MM B}}{\text{MM A}}}$
explanation	the sum of the pressures of the individual gases in a mixture equals the total pressure exerted by the mixture	amount of a gas in mixture is proportionate to the amount of its partial pressure	rate of gas A compared to the rate of gas B is equal to the square root of the inverse of their molar masses
when to use it	mixture of gases; only pressures given	mixture of gases; moles & total pressure given	when any form of the word "effusion" or "diffusion" is in the problem
specific units req'd?	any - but all values must have same unit of pressure	any - but all values must have same unit of pressure	no

Law	Molar Mass (from Ideal Gas Equation)	Density (from Ideal Gas Equation)
equation	$MM = \frac{g R T}{P V}$	$D = \frac{P MM}{R T}$
explanation	one gas at one set of conditions, asking about molar mass (no information about density given)	one gas at one set of conditions (involves density)
when to use it	when the problem gives: mass (grams), P, V, T	when density is mentioned (given or asked for in problem)
specific units req'd?	pressure = atm, volume = liters, temperature = Kelvins, molar mass = g/mole	pressure = atm, volume = liters, temperature = Kelvins, molar mass = g/mole, density = g/L