

**Types of Mixtures Notes**

\*What is the Tyndall Effect?

|            | Homogeneous or Heterogeneous | # of visible phases | Settling? | Filterable? | Tyndall Effect? * | Particle Size | Examples |
|------------|------------------------------|---------------------|-----------|-------------|-------------------|---------------|----------|
| Suspension |                              |                     |           |             |                   |               |          |
| Colloid    |                              |                     |           |             |                   |               |          |
| Solution   |                              |                     |           |             |                   |               |          |

**COMPONENTS OF A SOLUTION**

~ Solute:

~ Solvent:

\* What substance is the most common solvent?

\* What is it referred to as?

**TYPES OF SOLUTIONS**

~ Examples (solute/solvent)  
 solid/liquid  
 liquid/liquid  
 gas/liquid  
 solid/solid

**FACTORS AFFECTING THE RATE OF SOLUTION**

- 1.
- 2.
- 3.

**SOLUBILITY:**


~ Saturated:

~ Unsaturated:

~ Supersaturated:

\* How can you make a supersaturated solution?

**SOLUBILITY CURVES WORKSHEET**



- Which compound is *least* soluble at: (A) 20°C? (B) 80°C?
- Which substance is the *most* soluble at: (A) 10°C? (B) 50°C? (C) 90 °C?
- The solubility of which substance is *most* affected by changes in temperature?
- The solubility of which substance is *least* affected by changes in temperature?
- Are the following solutions saturated, unsaturated, or supersaturated?  
(Assume all are dissolved in 100 grams of water.)
  - 50 grams of NH<sub>4</sub>Cl at 50°C
  - 100 grams of NaNO<sub>3</sub> at 80°C
  - 30 grams of KNO<sub>3</sub> at 25°C
  - 51 grams of KCl at 80°C
  - 65 grams of NH<sub>4</sub>Cl at 70°C
  - 30 grams of NH<sub>3</sub> at 50°C
  - 10 grams of KClO<sub>3</sub> at 20°C

- 6.)  $\text{NH}_3$  is a gas. Describe what happens to its solubility as the temperature goes from  $20^\circ\text{C}$  to  $80^\circ\text{C}$ .
- 7.) Which two substances have the same solubility at  $68^\circ\text{C}$ ? What is the solubility?
- 8.) Which two substances have the same solubility at  $94^\circ\text{C}$ ? What is the solubility?
- 9.) For each of the following, indicate the temperature at which the solution described would be saturated. (Assume all are dissolved in 100 grams of water.)
- (A) 30 grams of  $\text{NH}_4\text{Cl}$
  - (B) 130 grams of  $\text{NaNO}_3$
  - (C) 50 grams of  $\text{Na}_2\text{SO}_4$
  - (D) 20 grams of  $\text{KNO}_3$
  - (E) 40 grams of  $\text{KCl}$
  - (F) 60 grams of  $\text{NH}_3$
- 10.) For each of these, indicate how many grams of solute (per 100 grams of water) will dissolve.
- (A)  $\text{NaNO}_3$  at  $70^\circ\text{C}$
  - (B)  $\text{NH}_4\text{Cl}$  at  $50^\circ\text{C}$
  - (C)  $\text{KI}$  at  $20^\circ\text{C}$
  - (D)  $\text{KClO}_3$  at  $90^\circ\text{C}$
- 11.) At  $40^\circ\text{C}$ , how many grams of  $\text{NaNO}_3$  will make a saturated solution if the  $\text{NaNO}_3$  is added to 100 grams of water?
- 12.) At  $80^\circ\text{C}$ , how many grams of  $\text{KCl}$  can be dissolved in 200 grams of water?
- 13.) At what temperature will 10 grams of  $\text{NH}_3$  dissolve completely in 100 grams of water to make a saturated solution?
- 14.) At  $40^\circ\text{C}$ , how many grams of  $\text{KNO}_3$  can be dissolved in 300 grams of water?
- 15.) At  $55^\circ\text{C}$ , how many grams of  $\text{NaNO}_3$  can be dissolved in 50 grams of water?
- 16.) At  $80^\circ\text{C}$ , you have a saturated solution of  $\text{KClO}_3$ . How many grams of solid precipitate will form if the solution is cooled to  $50^\circ\text{C}$ ?
- 17.) How many grams of  $\text{NaNO}_3$  precipitate will form if a saturated solution at  $70^\circ\text{C}$  is cooled to  $10^\circ\text{C}$ ?
- 18.) A solution contains 20 g of  $\text{NH}_4\text{Cl}$  at  $50^\circ\text{C}$ . How many more grams of  $\text{NH}_4\text{Cl}$  need to be added to the 100 grams of water for the solution to be saturated?

**FACTORS AFFECTING SOLUBILITY NOTES**

~ TEMPERATURE:

In a solid/liquid solution, solubility \_\_\_\_\_ as the temp. of the solvent increases.

In a gas/liquid solution, solubility \_\_\_\_\_ as the temp. of the solvent increases.

~ PRESSURE:

In a gas/liquid solution, solubility \_\_\_\_\_ as the pressure of the gas over the liquid decreases.

**GENERAL SOLUBILITY RULE: "LIKE DISSOLVES LIKE"**

- ~ Polar solvents / polar solutes:
- ~ Polar solvents / nonpolar solutes:
- ~ Nonpolar solvents / polar solutes:
- ~ Nonpolar solvents / nonpolar solutes:

**CONCENTRATION OF SOLUTIONS**

- ~ MOLARITY (M) =  $\frac{\text{moles of solute}}{\text{liters of solution}}$

EXAMPLE: How many liters of a 0.25 M solution can be made from 50.0 grams of  $\text{CaCl}_2$ ?

=====

~ DILUTIONS:  $M_1 V_1 = M_2 V_2$

EXAMPLE: What is the molarity of a solution made by diluting 37.5 mL of 12.0 M HCl to 150.0 mL?

=====

**Molarity & Dilutions Worksheet**

- 1.) What is the molarity of a solution in which 58.5 grams of NaCl are dissolved in 1.0 L of solution?
- 2.) What is the molarity of a solution in which 10.0 grams of  $\text{AgNO}_3$  is dissolved in 500. mL of solution?
- 3.) How many grams of  $\text{KNO}_3$  should be used to prepare 2.00 L or a 0.500 M solution?
- 4.) What volume of a 0.25 M solution could be made from 5.0 grams of KCl?
- 5.) How many grams of  $\text{CuSO}_4$  are needed to prepare 100. mL of a 0.10 M solution?
- 6.) How much 18 M sulfuric acid ( $\text{H}_2\text{SO}_4$ ) is needed to prepare 250 mL of a 6.0 M solution?
- 7.) 17 mL of 12 M hydrochloric acid (HCl) is diluted to 100. mL. What is the concentration of the new solution?
- 8.) To what volume should 25 mL of 15 M nitric acid ( $\text{HNO}_3$ ) be diluted to prepare a 3.0 M solution?
- 9.) To what volume should 50. mL of 12 M hydrochloric acid be diluted to produce a 4.0 M solution?
- 10.) If 25.0 mL of 18 M sulfuric acid is diluted to 550. mL, what is the concentration of the diluted solution?

Answers: 1.) 1.0 M; 2.) 0.118 M; 3.) 101 g; 4.) 0.27 L or 270 mL; 5.) 1.6 g; 6.) 83 mL; 7.) 2.0 M; 8.) 130 mL; 9.) 150 mL; 10.) 0.82 M



SPECTATOR IONS do not take part in a chemical reaction. They appear exactly the same on both sides of the equation. The spectator ions are the ones that get crossed out. They are simply "spectators" watching the chemical reaction take place – not taking part in it.

The spectator ions in the example are  $K^{+1} (aq)$  and  $NO_3^{-1} (aq)$ .

The "shortcut" for writing net ionic equations:

Step 1 – Determine the solubility of the products of the reaction.

Step 2 – Write the product of the net ionic equation. (The product of a net ionic equation will always be the insoluble/solid product.)

Step 3 – Determine the reactants using the product as a guide. (The reactants will be the positive ion of the insoluble/solid product written with its oxidation number and "(aq)" after it, and then the negative ion of the insoluble/solid product written with its oxidation number and "(aq)" after it.)

Step 4 – Balance the equation using coefficients, if necessary.

The spectator ions in a reaction will be the positive and negative ions of the soluble/aqueous product.

Example:  $KOH (aq) + Zn(NO_3)_2 (aq) \rightarrow KNO_3 ( ) + Zn(OH)_2 ( )$

Net ionic equation:

Spectator ions:

### **NET IONIC EQUATIONS WORKSHEET**

Determine the solubility of the products ((aq) or (s)). Then, write the net ionic equation. Also indicate the spectator ions for each equation.

1.)  $Na_2S (aq) + AgC_2H_3O_2 (aq) \rightarrow Ag_2S ( ) + NaC_2H_3O_2 ( )$

net ionic eqn:

spectator ions:

2.)  $Zn(NO_3)_2 (aq) + (NH_4)_2S (aq) \rightarrow NH_4NO_3 ( ) + ZnS ( )$

net ionic eqn:

spectator ions:

3.)  $K_2SO_4 (aq) + Ba(NO_3)_2 (aq) \rightarrow BaSO_4 ( ) + KNO_3 ( )$

net ionic eqn:

spectator ions:

4.)  $AgNO_3 (aq) + NaI (aq) \rightarrow AgI ( ) + NaNO_3 ( )$


net ionic eqn:

spectator ions:

5.)  $HgCl_2 (aq) + K_2S (aq) \rightarrow KCl ( ) + HgS ( )$


net ionic eqn:

spectator ions:




net ionic eqn:

spectator ions:




net ionic eqn:

spectator ions:




net ionic eqn:

spectator ions:



net ionic eqn:

spectator ions:



net ionic eqn:

spectator ions:

**UNIT 12 REVIEW WORKSHEET**

Part 1 – Solubility Curves - USE YOUR SOLUBILITY CURVE GRAPH TO ANSWER #1-4.

1. At what temperature does 135 grams of KI dissolved in 100 grams of water form a saturated solution?
2. How many grams of  $\text{KNO}_3$  will dissolve in 400 grams of water at  $60^\circ\text{C}$ ?
3. If 10 grams of  $\text{KClO}_3$  are dissolved in 100 grams of water at  $30^\circ\text{C}$ , is the solution saturated, unsaturated, or supersaturated?
4. How many grams of solid precipitate will form if a saturated  $\text{NaNO}_3$  solution is cooled from  $80^\circ\text{C}$  to  $20^\circ\text{C}$ ?

Part 2 – Concentration of Solutions

5. How many grams of  $\text{Fe}(\text{C}_2\text{H}_3\text{O}_2)_3$  are needed to dissolve to make 350. mL of a 2.50 M solution?
6. What is the molarity of a solution made by diluting 17.3 mL of 12 M hydrochloric acid to 550. mL?

Part 3 – Solubility – Write (s) or (aq) next to each product. Then, write the net ionic equation.

