


BRONSTED - LOWRY ACIDS & BASES WORKSHEET

According to Bronsted-Lowry theory, an acid is a proton (H^{+1}) donor, and a base is a proton acceptor.

Label the Bronsted-Lowry acids (A), bases (B), conjugate acids (CA), and conjugate bases (CB) in the following reactions.



- $C_6H_5NH_2 + H_2O \leftrightarrow C_6H_5NH_3^{+1} + OH^{-1}$
- $H_2SO_4 + OH^{-1} \leftrightarrow HSO_4^{-1} + H_2O$
- $HSO_4^{-1} + H_2O \leftrightarrow SO_4^{-2} + H_3O^{+1}$
- $HBr + OH^{-1} \leftrightarrow H_2O + Br^{-1}$
- $NH_3 + H_2O \leftrightarrow NH_4^{+1} + OH^{-1}$

CONJUGATE ACID-BASE PAIRS WORKSHEET

A conjugate base is what is left after an acid gives up its proton.
 A conjugate acid is what is made once a base gains a proton.

ACID	BASE
	HSO_4^{-1}
H_3PO_4	
	F^{-1}
	NO_3^{-1}
$H_2PO_4^{-1}$	
H_2O	
	SO_4^{-2}
HPO_4^{-2}	
NH_4^{+1}	

pH and pOH WORKSHEET

Part 1 - Calculate the pH and identify as acidic, basic, or neutral.

- $[H^{+1}] = 1.0 \times 10^{-6} M$ pH = _____ A B N
- $[H^{+1}] = 2.61 \times 10^{-2} M$ pH = _____ A B N
- $[H^{+1}] = 4.0 \times 10^{-9} M$ pH = _____ A B N
- $[H^{+1}] = 5.9 \times 10^{-12} M$ pH = _____ A B N
- $[H^{+1}] = 1.0 \times 10^{-7} M$ pH = _____ A B N

Part 2 - Calculate the $[H^{+1}]$ and identify as acidic, basic, or neutral.

6.) pH = 4.00 $[H^{+1}] = \underline{\hspace{2cm}}$ M A B N

7.) pH = 5.89 $[H^{+1}] = \underline{\hspace{2cm}}$ M A B N

8.) pH = 7.00 $[H^{+1}] = \underline{\hspace{2cm}}$ M A B N

9.) pH = 12.25 $[H^{+1}] = \underline{\hspace{2cm}}$ M A B N

10.) pH = 9.11 $[H^{+1}] = \underline{\hspace{2cm}}$ M A B N

Part 3 - Calculate the missing $[H^{+1}]$ or $[OH^{-1}]$ and identify as acidic, basic, or neutral.

11.) $[H^{+1}] = 4.2 \times 10^{-6}$ M $[OH^{-1}] = \underline{\hspace{2cm}}$ M A B N

12.) $[H^{+1}] = \underline{\hspace{2cm}}$ M $[OH^{-1}] = 4.3 \times 10^{-5}$ M A B N

Part 4 - Complete the following chart.

Solution	pH	pOH	$[H^{+1}]$	$[OH^{-1}]$
A	3.68			
B				8.60×10^{-5} M
C			1.80×10^{-9} M	
D		5.48		
E	10.84			
F			3.82×10^{-11} M	
G		2.85		

ACID-BASE TITRATION WORKSHEET

- 1.) A 25.0 mL sample of HCl was titrated to the endpoint with 15.0 mL of 2.0 M NaOH. What is the molarity of the HCl?
- 2.) A 10.0 mL sample of H_2SO_4 was exactly neutralized by 13.5 mL of 1.0 M KOH. What is the molarity of the H_2SO_4 ?
- 3.) How much 1.5 M NaOH is necessary to exactly neutralize 20.0 mL of 2.5 M H_3PO_4 ?
- 4.) How much of 0.50 M HNO_3 is necessary to titrate 25.0 mL of 0.050 M $Ca(OH)_2$ solution to the endpoint?
- 5.) What is the molarity of NaOH solution if 15.0 mL is exactly neutralized by 7.5 mL of a 0.020 M $HClO_3$ solution?

32. What is the concentration (molarity) of NaOH if 15.3 mL are needed to completely neutralize 20.4 mL of 2.50 M H₂SO₄?
33. What volume of 1.50 M Ca(OH)₂ is needed to reach the endpoint of a titration using 17.2 mL of 3.00 M H₃PO₄?

Part 6 – Fill in the following chart.

pOH	pH	[OH ⁻¹]	[H ⁺¹]	A, B, or N
4.63				
	11.75			
		2.96 x 10 ⁻⁸ M		
			5.27 x 10 ⁻⁹ M	

ACID – BASE TITRATION LAB

PURPOSE: - to determine the molarity of an aqueous solution of hydrochloric acid (HCl)
- to learn and practice the technique for titration

MATERIALS:


EQUIPMENT

- 2 burets (labeled "A" and "B")
- double buret clamp
- ring stand
- Erlenmeyer flask

CHEMICALS

- _____ M NaOH
- phenolphthalein
- ? M HCl

READING A BURET:



PROCEDURE:

- 1.) Obtain approximately 10 mL of HCl from the buret labeled "A" for acid in the Erlenmeyer flask. (It is not necessary to have EXACTLY 10.00 mL. However, the volume of the HCl should be measured to the nearest hundredth.)
- 2.) Add 3 or 4 drops of phenolphthalein. (Only a few drops are needed. Adding more than 3 or 4 drops will not make the reaction any bigger or better.)
- 3.) Add NaOH from the buret labeled "B" for base SLOWLY until the solution changes color. (When the solution changes color, the acid is neutralized. Ideally, the solution in the flask should be a very faint shade of the color.)

DATA TABLE:

Unit 14 – Acids & Bases

		TRIAL 1	TRIAL 2	TRIAL 3
1	Reading of HCl (at start)	mL	mL	mL
2	Reading of HCl (at end)	mL	mL	mL
3	Volume of HCl used	mL	mL	mL
4	Reading of NaOH (at start)	mL	mL	mL
5	Reading of NaOH (at end)	mL	mL	mL
6	Volume of NaOH used	mL	mL	mL

QUESTIONS AND CALCULATIONS:

- 1.) Write a balanced equation for the neutralization reaction that occurs between hydrochloric acid (HCl) and sodium hydroxide (NaOH).
- 2.) What is the mole ratio between HCl and NaOH in the balanced equation?
- 3.) What volume of HCl was used in TRIAL 1? Show how you determined this volume.
- 4.) What volume of NaOH was used in TRIAL 1? Show how you determined this volume.
- 5.) The NaOH had a concentration of _____ M. Using your data, what is the concentration (molarity) of the hydrochloric acid? (CLEARLY and LEGIBLY show your calculations in detail!)
- 6.) If your group was able to do more than one trial, what was the average molarity?