


Electrons, Energy, & the Electromagnetic Spectrum Notes

Simplified, 2-D Bohr Model:

Figure 1



orbits, paths, shells, rings = ENERGY LEVELS


As the name "energy level" implies, there is a specific amount of energy associated with each energy level.

Electrons are lazy - they will occupy the location that requires the least amount of energy. This is called the AUFBAU principle.

lowest energy state = GROUND STATE

Figure 2


For hydrogen in the ground state...



- ~ Electron is located in energy level 1 (E1) because E1 is closest to the nucleus. Nucleus is positively-charged & attracts negatively-charged electrons.
- ~ Therefore, low amount of energy needed to be in E1.

Figure 3


If energy (from an outside source) is added to the atom...



Electrons jump to a higher energy level because energy is absorbed. (Electron is now in the EXCITED STATE.)

As soon as the electron jumps to a higher level, it immediately falls back to a lower energy level. When the electron falls, energy is released.

The energy is released as electromagnetic radiation. The type of electromagnetic radiation produced depends on the difference in energy between the energy levels.



Light Calculation Notes

Here's how the type/form of EM radiation can be determined...

The amount of energy released when an electron falls from a higher to a lower energy level is directly proportional to its frequency.

The calculation follows the equation: $E = h \cdot \nu$

E = Energy (unit is J)

h = Planck's Constant (6.626×10^{-34} Js)

ν = frequency (unit is Hz or $1/\text{second}$)

EXAMPLE 1: A particle of EM radiation has an energy of 1.15×10^{-16} J. What is its frequency?

$$1.15 \times 10^{-16} \text{ J} = 6.626 \times 10^{-34} \text{ Js} \cdot \nu$$

$$\nu = 1.74 \times 10^{17} \text{ Hz}$$

The type of electromagnetic radiation can be determined if one knows the wavelength. The wavelength is inversely proportional to the frequency.

The calculation follows the equation: $c = \lambda \cdot \nu$

c = speed of light (3.00×10^8 m/s)

λ = wavelength (unit is m)

ν = frequency (unit is Hz or $1/\text{s}$)

EXAMPLE 2: What is the wavelength of the same particle from EXAMPLE 1?

$$3.00 \times 10^8 \text{ m/s} = \lambda \cdot 1.74 \times 10^{17} \text{ Hz}$$

$$\lambda = 1.72 \times 10^{-9} \text{ m}$$

EXAMPLE 3: What type of electromagnetic radiation is the particle from EXAMPLE 1?

answer for wavelength is 10^{-9} so use chart below to determine...

x-rays or ultraviolet (either one is acceptable)


PROBLEMS FOR YOU TO TRY ON YOUR OWN...

1.) A particle of EM radiation has a frequency of 4.80×10^{14} Hz.

(A) How much energy does this particle have?

- (B) What is the wavelength of this particle?
 (C) What specific type of electromagnetic radiation does this particle represent?
- 2.) A particle of electromagnetic radiation has 2.39×10^{-13} Joules of energy.
 (A) What is the wavelength of this particle?
 (B) What type of electromagnetic radiation does this particle represent?

Electromagnetic Spectrum
 (measurement in meters)



EM SPECTRUM, WAVELENGTH, FREQUENCY, AND ENERGY WORKSHEET

- 1.) Look at the EM spectrum below to answer this question.
 As you move across the visible light spectrum from red to violet...
 (A) Does the wavelength increase or decrease?
 (B) Does the frequency increase or decrease?
 (C) Does the energy increase or decrease?
- 2.) A beam of microwaves has a frequency of 1.0×10^9 Hz. A radar beam has a frequency of 5.0×10^{11} Hz.
 Which type (microwave or radar)...
 (A) has a longer wavelength?
 (B) is closer to visible light on the EM spectrum?
 (C) is closer to x-rays in frequency value?
- 3.) What is the frequency of an EM radiation wave if its wavelength is 3.6×10^{-9} meters?
- 4.) A beam of EM radiation has a wavelength of 4.257×10^{-7} cm. What is its frequency?
- 5.) A photon of light has a wavelength of 3.20×10^5 meters. Find...
 (A) the frequency
 (B) the energy
 (C) the region of the EM spectrum/type of radiation
- 6.) A photon has an energy of 4.00×10^{-19} J. Find...
 (A) the frequency
 (B) the wavelength
 (C) the region of the EM spectrum/type of radiation
- 7.) A bright line spectrum contains a line with a wavelength of 518 nm. Determine...
 (A) the wavelength in meters
 (B) the frequency
 (C) the energy
 (D) the color

*8.) Cobalt-60 is an artificial radioisotope that is produced in a nuclear reactor for use as a gamma ray source in the treatment of certain types of cancer. If the wavelength of the gamma radiation from a cobalt-60 source is 1.00×10^{-3} nm, calculate the energy of a photon of this radiation.

Light Calculations Notes:

- * Frequency and wavelength are _____ proportional
- * Energy and frequency are _____ proportional

Light as a Particle Notes:

- * Object emits energy in small, specific amounts (called _____)
- * _____: particle of EM radiation carrying a quantum of energy
- * Einstein suggested that light had properties of both waves and particles
Referred to as the _____ of light

Quantum Theory Notes:

- * When atom falls from excited state to ground state, _____
- * Energy of photon = difference _____
- * Energy states of atoms are fixed

Bohr model of the hydrogen atom Notes:

- * said that e- circled the nucleus in fixed paths
- * when in path, has fixed amount of energy
- * e- cannot exist in space between path
- * drawback of Bohr's model =

Example Questions:

1. Which has a longer wavelength - microwaves or x-rays?
2. Which has higher frequency - radio waves or ultraviolet?
3. Which has more energy - gamma rays or visible light?

EMISSION & ABSORPTION SPECTRA NOTES

According to the Bohr atomic model, electrons orbit the nucleus within specific energy levels. These levels are defined by unique amounts of energy. Electrons possessing the lowest energy are found in the levels closest to the nucleus. Electrons with higher energy are located in progressively more distant energy levels.

If an electron absorbs sufficient energy to bridge the "gap" between energy levels, the electron may jump to a higher level. Since this change results in a vacant lower orbital, this configuration is unstable. The "excited" electron releases its newly acquired energy as it falls back to its initial or ground state. Often, the excited electrons acquire sufficient energy to make several energy level transitions. When these electrons return to the ground state, several distinct energy emissions occur. The energy that the electrons absorb is often of a thermal or electrical nature, and the energy that an electron emits when returning to the ground state is called electromagnetic radiation.

In 1900, Max Planck studied visible emissions from hot glowing solids. He proposed that light was emitted in "packets" of energy called quanta, and that the energy of each packet was proportional to the frequency of the light wave. According to Einstein and Planck, the energy of the packet could be expressed as the product of the frequency (ν) of emitted light and Planck's constant (h). The equation is written as $E = h \cdot \nu$

If white light passes through a prism or diffraction grating, its component wavelengths are bent at different angles. This process produces a rainbow of distinct colors known as a continuous spectrum. If, however, the light emitted from hot gases or energized ions is viewed in a similar manner, isolated bands of color are observed. These bands form characteristic patterns - unique to each element. They are known as bright line spectra or emission spectra.

By analyzing the emission spectrum of hydrogen gas, Bohr was able to calculate the energy content of the major electron levels. Although the electron structure as suggested by his planetary model has been modified according to modern quantum theory, his description and analysis of spectral emission lines are still valid.

In addition to the fundamental role of spectroscopy played in the development of today's atomic model, this technique can also be used in the identification of elements. Since the atoms of each element contain unique arrangements of electrons, emission lines can be used as spectral fingerprints. Even without a spectroscope, this type of identification is possible since the major spectral lines will alter the color.

PROPERTIES OF LIGHT WORKSHEET

Part 1 - Select the best answer

1. Which has a longer wavelength, orange or violet light?
2. Which has a higher energy, x-rays or gamma rays?
3. Which has a lower frequency, radio waves or green light?
4. Which has the shortest wavelength, violet or ultraviolet light?
5. Which has lower energy, infrared light or x-rays?

Part 2 - Fill in the blanks

6. _____ formed a theory to explain the structure of an atom by revising physical theories.
7. As the energy level increases, the amount of energy an electron will possess _____.
8. Electrons give off energy in finite amounts called _____ when returning to the ground state.
9. When this energy is released in the form of light it is called a _____.
10. The speed of light = _____ (give number and units)
11. The symbol for wavelength is _____.
12. In the equation $c = \lambda \cdot \nu$, c represents _____, ν represents _____, and λ represents _____.
13. In the equation $c = \lambda \cdot \nu$, λ and ν are _____ proportional.
14. In the equation $E = h \cdot \nu$, h represents _____ and E represents _____.
15. In the equation $E = h \cdot \nu$, E and ν are _____ proportional.
16. Bohr chose the element _____ to prove his theory.


Part 3 - True or False

17. Electrons may regularly occupy spaces between energy levels.
18. The varying wavelengths on the electromagnetic radiation spectrum travel at different speeds.
19. Atoms release energy when electrons jump to higher energy levels.

ELECTRON ARRANGEMENT NOTES

Heisenberg Uncertainty Principle:

GENERAL LOCATION -----



SPECIFIC LOCATION -----

ENERGY LEVELS


- divisions of the electron cloud
- numbered consecutively from closest to farthest away from nucleus



Electrons will occupy the location with the lowest amount of energy.


SUBLEVELS

- divisions of energy levels
- designated by letters (s, p, d, f)
- number of sublevels in an energy level = # of the energy



ORBITALS

- divisions of sublevels
- number of orbitals in an energy level =
- "s" sublevel has 1 orbital; "p" sublevel has 3 orbitals; "d" sublevel has 5 orbitals; "f" sublevel has 7 orbitals



HOW MANY ELECTRONS CAN AN ORBITAL HOLD?

HOW MANY ELECTRONS CAN EACH SUBLEVEL HOLD?

"s" = e⁻

"p" = e⁻




"d" = e⁻

"f" = e⁻

11. Sublevels are divided into _____.
12. Each orbital can hold up to _____ electrons.
13. Sketch the shapes of the orbitals for the sublevels listed.
 (A) s: (B) p_x: (C) p_y: (D) p_z:
14. How many orbitals are in each sublevel?
 (A) s _____ (B) p _____ (C) d _____ (D) f _____

More Electron Arrangement NOTES




Examples:



HOEL (Highest Occupied Energy Level): energy level furthest from the nucleus that contains at least one electron

How to determine this using electron configuration?



~ largest non-exponent number



Valence Electrons: electrons in the HOEL

How to determine this using electron configuration?

~ add up exponents of terms in HOEL






Noble Gas Configuration: shortcut for electron configuration

How is it written?

~ [symbol for noble gas closest to element with lower atomic #]


~ [after brackets] write the rest of the configuration as if you had written the configuration for the noble gas

NOTE: ending of electron configuration and noble gas configuration should be the same



Orbital Notation: drawing of how electrons are arranged in orbitals; will only need to do this for the HOEL

*NOTE: $\underline{\quad}$ = orbital \uparrow or \downarrow = electrons



Dot Diagrams: symbol represents nucleus and non-valence ("inner-shell") electrons; dots around symbol represent valence electrons




QUANTUM NUMBERS NOTES

- ~ describe one specific electron
- ~ 1st quantum number = PRINCIPAL QUANTUM NUMBER
 - ~ abbreviated "n"
 - ~ tells the energy level the electron is located in
 - ~ n = number of the energy level
 - ~ 1st energy level: n = 1, 4th energy level: n = 4, etc.
- ~ 2nd quantum number = ANGULAR MOMENTUM QUANTUM NUMBER
 - ~ abbreviated "l"
 - ~ tells the sublevel the electron is located in
 - ~ tells shape of orbital
 - ~ "s" sublevel: l = 0, "p" sublevel: l = 1, "d" sublevel: l = 2, "f" sublevel: l = 3
- ~ 3rd quantum number = MAGNETIC QUANTUM NUMBER
 - ~ abbreviated "m"
 - ~ tells which orbital the electron is in
 - ~ tells orientation of orbital around nucleus
 - ~ m = - l .. + l
- ~ 4th quantum number = SPIN QUANTUM NUMBER
 - ~ abbreviated "s"
 - ~ tells which electron is being described
 - ~ tells which way electron is spinning
 - ~ s = -1/2 or +1/2

Pauli Exclusion Principle:

EXAMPLE QUESTIONS:

1.) What are the 4 quantum numbers for the following electron?



2.) If the electron in question 1 was the last electron added, what element would it be?


3.) Draw in the electron (and the orbital notation) for the electron with the following quantum numbers.

$$n = 3 \quad \ell = 2 \quad m = -1 \quad s = -\frac{1}{2}$$

4.) How many electrons in an atom can have the quantum numbers $n = 3$ and $\ell = 1$?

5.) What are the four quantum numbers for the electron circled in the diagram below?

$$n = \quad \ell = \quad m = \quad s =$$



QUANTUM NUMBERS WORKSHEET

	Element	1s	2s	2p	3s	3p	3d	4s	4p	4d	4f	5s
1.)	K	-	-	---	-	↑	---	-	---	---	---	-
2.)	O	-	↑	---	-	---	---	-	---	---	---	-
3.)	Ar	-	-	↓	-	---	---	-	---	---	---	-
4.)	Br	-	-	---	-	---	---	-	---	---	---	-
5.)	Rb	-	-	---	-	---	---	-	---	---	---	-
6.)	Co	-	-	---	-	---	---	-	---	---	---	-
7.)	Se	-	-	---	-	---	---	-	---	---	---	-
8.)	B	-	-	---	-	---	---	-	---	---	---	-
9.)	P	-	-	---	-	---	---	-	---	---	---	-

** Each question below corresponds to the number in the table. **

For 1 - 3, give the four quantum numbers for the electron indicated.

1.) $n =$ $\ell =$ $m =$ $s =$

2.) $n =$ $\ell =$ $m =$ $s =$

3.) $n =$ $\ell =$ $m =$ $s =$

For 4 - 9, draw in the electron with the following sets of quantum numbers.

4.) $n = 3$ $\ell = 2$ $m = 0$ $s = +\frac{1}{2}$ 5.) $n = 3$ $\ell = 1$ $m = +2$ $s = +\frac{1}{2}$

6.) $n = 4$ $\ell = 1$ $m = +1$ $s = -\frac{1}{2}$ 7.) $n = 2$ $\ell = 1$ $m = 0$ $s = +\frac{1}{2}$

8.) $n = 3$ $\ell = 2$ $m = -1$ $s = -\frac{1}{2}$ 9.) $n = 2$ $\ell = 1$ $m = -1$ $s = -\frac{1}{2}$

10.) Does an electron with this set of quantum numbers exist in the element calcium?

$$n = 4 \quad \ell = 1 \quad m = 0 \quad s = -\frac{1}{2}$$

Determine the element whose outermost electron (last electron added) is being defined by the following quantum numbers.

11.) $n = 1$ $\ell = 0$ $m = 0$ $s = -\frac{1}{2}$

12.) $n = 4$ $\ell = 1$ $m = 1$ $s = -\frac{1}{2}$

13.) $n = 3$ $\ell = 1$ $m = -1$ $s = +\frac{1}{2}$

14.) $n = 4$ $\ell = 0$ $m = 0$ $s = +\frac{1}{2}$

15.) $n = 3$ $\ell = 2$ $m = -2$ $s = -\frac{1}{2}$

Unit 4 Review Worksheet

Section I - Problems Given: $E = h \cdot \nu$ $h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$
 $c = \lambda \cdot \nu$ $c = 3.00 \times 10^8 \text{ m/s}$

1. What is the frequency of a wave with a wavelength of 3.5×10^{-4} m?
2. What is the energy of a photon with a frequency of 5.41×10^{17} Hz?
3. What type of electromagnetic radiation is described in question 2?

Section II - Electromagnetic Spectrum

4. Label both ends of the spectrum with high/low frequency, high/low energy, and long/short wavelength
radio waves microwaves infrared light ROYGBIV ultraviolet light x-rays gamma rays
5. Which has a higher energy, gamma or x-rays?
6. Which has a shorter wavelength, radio or ultraviolet?
7. Which has a lower frequency, yellow or green light?
8. In the equation $E = h \cdot \nu$, energy and frequency are _____ proportional.
9. In the equation $c = \lambda \cdot \nu$, wavelength and frequency are _____ proportional.
10. The symbol for wavelength is _____.
11. Electrons give off energy in the form of a _____ when returning to the ground state.
12. Which scientist proposed the idea that electrons travel around the nucleus in fixed paths?
13. When an electron moves from the ground state to the excited state, energy is _____.
14. Bohr chose the element _____ to prove his theory.
15. The dual wave-particle nature of electrons describes how the electrons in atoms can behave as _____ and _____.

Section III - Electrons

16. What is an electron cloud?
17. Who proposed the uncertainty principle?
18. Who is credited with the idea that electrons are placed in the lowest energy level first?
19. What rule requires that each of the "p" orbitals (at a particular energy level) receive one electron before any of the orbitals can have two electrons?
20. What is the maximum number of electrons in any orbital?
21. The principal quantum number, n, indicates the _____.
22. The maximum number of electrons in an energy level can be determined by the equation _____
 That means the maximum number of electrons in the 3rd energy level is _____.
23. The number of sublevels in any energy level can be determined by _____.
24. The number of orbitals in an energy level can be determined by the equation _____.
 So, the 3rd energy level has _____ orbitals. (_____ is/are "s" orbitals, _____ is/are "p" orbitals, and _____ is/are "d" orbitals)
25. List the four sublevels according to increasing energy.
26. The "s" sublevel is shaped like a _____ and has _____ orbitals.
27. A "p" sublevel is shaped like a _____ and has _____ orbitals.
28. The "d" sublevel has _____ orbitals and the "f" sublevel has _____ orbitals.

Section IV - Electron configuration, noble gas configuration, valence electrons, orbital notations

29. What is the electron configuration for phosphorus?
30. How many total electrons are in a neutral atom of phosphorus?
31. Write the noble gas configuration for phosphorus.
32. What is the highest occupied energy level for phosphorus?

33. What is the atomic number of phosphorus?
 34. Draw the orbital notation for phosphorus.
 35. Circle the last electron added to phosphorus. What are the four quantum numbers for this electron?
 n = ℓ = m = s =

36. How many electrons are in the highest occupied energy level of phosphorus?
 37. How many inner-shell electrons does phosphorus have?
 38. In which orbitals are the inner-shell electrons located?
 39. Draw the electron dot diagram for phosphorus.

Section V - Quantum numbers (Honors level only)

40. How many electrons can be described by the quantum numbers $n = 3$ and $ℓ = 1$?
 41. How many electrons in an atom can have the quantum numbers $n = 2$ and $ℓ = 3$?
 42. How many electrons can have the value $n = 3$?
 43. How many electrons in an atom have the quantum numbers $n = 4$ and $ℓ = 2$?
 44. Which of the following sets of quantum numbers does NOT represent a possible set of quantum numbers?
 (There may be more than one correct answer.)

	<u>n</u>	<u>ℓ</u>	<u>m</u>	<u>s</u>
(A)	4	8	-4	$1/2$
(B)	6	5	-5	$1/2$
(C)	3	2	2	$1/2$
(D)	6	0	1	$1/2$