

**LOCATING MAIN GROUP ELEMENTS ON THE PERIODIC TABLE NOTES**

Given the electron configuration or noble gas configuration for an element, it is possible to determine its location on the Periodic Table without actually looking at a Periodic Table.

\* To tell which period this element is in... ~ find the highest occupied energy level for this element  
 You can do this by... ~ finding the largest coefficient number

**The largest coefficient number is the number of the period where the element is located.**

\* To tell which "block" (s, p, d, f) this element is in... ~ find the highest occupied sublevel for this element  
 You can do this by... ~ finding the last lowercase letter written

**The last lowercase letter written in the configuration is the "block" where the element is located.**

\* To tell which group this element is in... ~ find the highest occupied energy level for this element  
 Then... ~ add up the exponents of the largest coefficient number

**This gives you the number of valence electrons in the element.**

You will then know that 1 valence e- indicates that the element is in Group 1, 2 valence e- indicates that the element is in Group 2, 3 valence e- indicates that the element is in Group 13, 4 valence e- indicates that the element is in Group 14, 5 valence e- indicates that the element is in Group 15, 6 valence e- indicates that the element is in Group 16, 7 valence e- indicates that the element is in Group 17, and 8 valence e- indicates that the element is in Group 18.

Look at the following EXAMPLE: [Ar] 4s<sup>2</sup> 3d<sup>10</sup> 4p<sup>5</sup>

It is possible to tell the period, group, and "block" where this element is located.

- \* Period -- largest coefficient number is 4, so element is in Period 4
- \* Block-- last lowercase letter written is "p", so element is in "p" block
- \* Group-- largest coefficient number is 4... 2 electrons in 4s, 5 electrons in 4p --> total of 7 valence electrons, so this element is in Group 17.

**LOCATION OF ELEMENTS WORKSHEET**

	Noble Gas Config.	Period	Block (s, p, d, f)	Group
1	[Ne] 3s <sup>2</sup> 3p <sup>2</sup>			
2	[Ar] 4s <sup>2</sup> 3d <sup>10</sup> 4p <sup>6</sup>			
3	[Xe] 6s <sup>2</sup>			
4	[Kr] 5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>5</sup>			
5	[Ar] 4s <sup>2</sup> 3d <sup>10</sup> 4p <sup>1</sup>			
6	[He] 2s <sup>2</sup> 2p <sup>3</sup>			
7	[Kr] 5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>4</sup>			
8	[He] 2s <sup>1</sup>			
9	[Xe] 6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>2</sup>			
10	[Rn] 7s <sup>2</sup>			

MAIN GROUP (GROUPS 1, 2, AND 13 - 18) ELEMENTS NOTES

GROUP	NAME	ENDING OF CONFIG.	# OF VALENCE e-	ELECTRON DOT DIAGRAM	WANTS TO (lose or gain) TO BE LIKE NOBLE GAS?	CHARGE OF ION & OXID-ATION #	ION (smaller or larger) THAN ATOM
1							
2							
13							
14							
15							
16							
17							
18							

HISTORY OF THE PERIODIC TABLE NOTES

I. Mendeleev and Chemical Periodicity

- A. Wanted to organize elements according to their \_\_\_\_\_
- B. When elements were arranged in order of increasing atomic mass\*, similarities in chemical properties appeared at regular intervals (\_\_\_\_\_)
- C. \*Several elements did not quite fit this pattern - Mendeleev put elements with similar \_\_\_\_\_ in the same column or group
- D. 1871 - Mendeleev predicted the existence and properties of several (then undiscovered) elements. These elements were:
- E. Within 15 years, those elements with those properties had been discovered

II. Moseley and the Periodic Law

- A. When elements were arranged in order of increasing \_\_\_\_\_, there was a distinct regular pattern.
- B. \_\_\_\_\_: The physical and chemical properties of the elements are periodic functions of their atomic numbers.

## UNIT 5 - PERIODIC TABLE & PERIODIC LAW

- C. In other words, when elements are arranged in order of increasing atomic number, elements with similar properties appear at regular intervals.
  - D. Bottom line = elements in the same group have similar properties
- III. Modern Periodic Table: arrangement of the elements in order of their atomic numbers so that elements with similar properties fall in the same group

### ELECTRON CONFIGURATION & THE PERIODIC TABLE NOTES

#### I. Stability of Noble Gases

- A. Noble gases undergo very few chemical reactions - why?
- B. Highest occupied energy level contains \_\_\_\_\_
- C. Electrons in the highest occupied energy level are what determines an element's \_\_\_\_\_

#### II. Periods and Blocks of the Periodic Table

- A. Horizontal row = \_\_\_\_\_; 7 on modern Periodic Table
- B. Length of period determined by the sublevels being filled in that period
- C. Period 1: only \_\_\_ sublevel being filled; can hold a maximum of \_\_\_ electrons; period contains \_\_\_ elements
- D. Period 4: \_\_\_, \_\_\_, and \_\_\_ sublevels being filled; s can hold \_\_\_ electrons, d can hold \_\_\_ electrons, & p can hold \_\_\_ electrons; total of \_\_\_ electrons; Period 4 contains \_\_\_ elements
- E. Period can be determined from the element's electron configuration
  - 1. Bromine:  $[\text{Ar}] 4s^2 3d^{10} 4p^5$
  - 2. Highest number in front of letter is the element's highest occupied \_\_\_\_\_ - tells which period the element is in
  - 3. For bromine, \_\_\_ is highest number, so it is in Period \_\_\_\_

#### III. The "s" block elements: Groups 1 and 2

- A. Group 1 - Alkali Metals
  - 1. generalized outermost energy level (valence) electron configuration:
  - 2. silvery appearance
  - 3. soft enough to cut with a knife
  - 4. not found in nature as free elements - they're always part of a compound
- B. Group 2 - Alkaline Earth Metals
  - 1. generalized valence electron configuration:
  - 2. harder, stronger, more dense than Group 1
  - 3. less reactive than Group 1, but still not found in nature as free elements
- C. Exceptions: Hydrogen and Helium
  - 1. Hydrogen (H)
    - a. electron configuration:
    - b. properties do not resemble those of any other element on the periodic table
  - 2. Helium (He)
    - a. electron configuration:
    - b. in Group 18 because

#### IV. The "d" block elements: Groups 3 - 12

- A. called
- B. have typical metallic properties: ductile, malleable, shiny, solid, conduct electricity
- C. less reactive than "s" block elements
- D. found in nature as free elements
- E. usual ending of electron configuration:

#### V. The "p" block elements: Grps 13 - 18 ("s" & "p" block elements together referred to as MAIN GROUP elements)

- A. ending electron configurations of \_\_\_\_\_ through \_\_\_\_\_
- B. properties vary greatly b/c there are metals, metalloids, and nonmetals
- C. Group 17 - Halogens
  - 1. most reactive nonmetals
  - 2. seven electrons in outermost energy level
- E. "p" block metals are harder and more dense than "s" block, but not as hard or dense as the "d" block metals

### PERIODIC TRENDS NOTES

Electronegativity/Electron Affinity (EN/EA): measure of how much an atom wants to gain an electron


**EN/EA Left to Right across a Period: INCREASES (not including Noble Gases)**

Why?

\* Elements on the left side of the P.T. (metals) want to lose electrons. Elements on the right side of the P.T. (nonmetals) want to gain electrons. Trend does not include Noble Gases because these elements do not want to lose or gain electrons.

**EN/EA Top to Bottom in a Group: DECREASES**

Why?



all Group 17 elements want to gain 1 electron to have a full "s" and "p" sublevel in their H.O.E.L.

Elements at the top have a stronger attraction for electrons (higher electronegativity) because their nucleus (where positive charges are) is physically closer to the element's H.O.E.L.

Additionally, more energy levels not only results in a larger distance between the nucleus and the H.O.E.L., but also there are more electrons in those energy levels.

These "inner-shell" electrons interfere with (and decrease) the strength of the hold the nucleus has for the valence electron(s).

\*

This interference (and resulting decreased "hold") is referred to as the SHIELDING EFFECT.

=====  
Ionization Energy (IE): amount of energy required to remove an atom's most loosely held electron


**IE Left to Right across a Period: INCREASES**

Why?

\* Elements on the left side of the P.T. (metals) want to lose electrons. Therefore, it will not require much energy to remove an electron. Elements on the right side of the P.T. (nonmetals) want to gain electrons. Consequently, a lot of energy will be needed to remove (take away) an electron.

**IE Top to Bottom in a Group: DECREASES**

Why?



Elements in Group 1 want to lose their 1 valence electron. Elements at the bottom of the P.T. lose their valence electron more easily (and with less required energy) than elements at the top of the P.T.

The reason why the elements at the bottom of the P.T. lose their electron easier is because of the decreased hold that the positive nucleus has for negative electrons - particularly the valence electron.


This decreased hold is due to the interference of the inner-shell electrons. The greater shielding effect (at the bottom of the P.T.) the lower the amount of energy needed to remove a valence electron.

\*

=====  
Atomic Radius (AR): distance from the nucleus to the H.O.E.L.

**AR Left to Right across a Period: DECREASES**

Why?



In all Period 2 elements, there are 2 occupied energy levels. As atomic number increases in a period (from left to right), the number of protons in the nucleus of each atom increases.

This increase in positive charges (protons in the nucleus) allows for a stronger attraction ("pull") for the negatively-charged electrons. Even though there are more electrons in each atom, the electrons are distributed over the same number of energy levels.


This stronger attraction allows for a greater pull by the nucleus. This results in the electrons being physically pulled closer to the nucleus. The result is a smaller distance between the nucleus and the H.O.E.L.

This attraction is referred to as EFFECTIVE NUCLEAR CHARGE.

**AR Top to Bottom in a Group: INCREASES**

Why?

\* There are more occupied energy levels as you move towards the bottom of the P.T.



Metallic Character: how easily an atom will lose valence electrons (easier to lose = more metallic = more reactive METAL)

Which metal loses its valence electron(s) most easily? Fr

Why?

\* Francium has one valence electron. It is more reactive than elements at the top of Group 1 because there are many inner shell electrons that decrease the attraction the nucleus has for the valence electrons.

**Nonmetallic Character:** how easily an atom will gain electrons (easier to gain = more nonmetallic = more reactive NONMETAL)

Which nonmetal gains electron(s) most easily? F

Why?

\* Fluorine has seven valence electrons. It is more reactive than elements at the bottom of Group 17 because there are only a few inner shell electrons. Consequently, the nucleus has a strong attraction for other electrons.

**PERIODIC TRENDS (multiple choice w/o using Periodic Table) WORKSHEET**

- 1.) Which element is most metallic? Group 14, Period \_\_\_  
(A) 2 (B) 3 (C) 4 (D) 5
- 2.) Which element is most nonmetallic? Group 16, Period \_\_\_  
(A) 2 (B) 3 (C) 4 (D) 5
- 3.) Which has the largest atomic radius? Group \_\_\_, Period 2  
(A) 1 (B) 13 (C) 15 (D) 17
- 4.) Which has the highest ionization energy? Group 2, Period \_\_\_  
(A) 3 (B) 4 (C) 5 (D) 6
- 5.) Which has the most metallic properties? Group \_\_\_, Period 5  
(A) 13 (B) 14 (C) 15 (D) 16
- 6.) Which has the greatest electron affinity?  
(A) Grp 16, Pd 4 (B) Grp 16, Pd 5 (C) Grp 17, Pd 5 (D) Grp 17, Pd 4
- 7.) Which has the smallest atomic radius? Group 15, Period \_\_\_  
(A) 2 (B) 3 (C) 4 (D) 5
- 8.) Which has the lowest electron affinity? Group \_\_\_, Period 3  
(A) 13 (B) 15 (C) 17 (D) 18
- 9.) Which has the lowest ionization energy? Group 1, Period \_\_\_  
(A) 2 (B) 3 (C) 4 (D) 5
- 10.) Which has the most metallic properties?  
(A) Grp 15, Pd 5 (B) Grp 16, Pd 5 (C) Grp 15, Pd 6 (D) Grp 16, Pd 6
- 11.) Which would most easily lose its valence electrons?  
(A) Grp 1, Pd 3 (B) Grp 14, Pd 2 (C) Grp 17, Pd 3 (D) Grp 18, Pd 2
- 12.) Which would most easily gain electrons?  
(A) Grp 13, Pd 3 (B) Grp 14, Pd 2 (C) Grp 15, Pd 2 (D) Grp 17, Pd 3
- 13.) Which has an octet of electrons in its outermost energy level?  
(A) Grp 13, Pd 3 (B) Grp 14, Pd 2 (C) Grp 18, Pd 2 (D) Grp 17, Pd 5
- 14.) Which has chemical properties most similar to [Ar] 4s<sup>1</sup>? Group \_\_\_, Period 3  
(A) 1 (B) 2 (C) 13 (D) 14
- 15.) Which is most reactive? Group \_\_\_, Period 2  
(A) 14 (B) 15 (C) 17 (D) 18
- 16.) Which is most reactive?  
(A) Grp 13, Pd 2 (B) Grp 1, Pd 5 (C) Grp 2, Pd 5 (D) Grp 13, Pd 5

- 17.) Which has chemical properties most similar to  $[\text{Ne}] 3s^2 3p^5$ ?  
 (A) Grp 16, Pd 3      (B) Grp 18, Pd 3      (C) Grp 17, Pd 4      (D) Grp 18, Pd 2
- 18.) Which would never be found in the free state?  
 (A) Grp 1, Pd 4      (B) Grp 13, Pd 3      (C) Grp 15, Pd 3      (D) Grp 14, Pd 4
- 19.) Which is the least reactive gas?      Group \_\_, Period 2  
 (A) 16      (B) 15      (C) 17      (D) 18
- 20.) Which is the most reactive gas?      Group \_\_, Period 2  
 (A) 16      (B) 15      (C) 17      (D) 18
- 21.) Which would never be in a compound?  
 (A) Grp 1, Pd 1      (B) Grp 18, Pd 1      (C) Grp 13, Pd 2      (D) Grp 1, Pd 2
- 22.) Which would be found in the "d" block of elements?  
 (A) Grp 1, Pd 3      (B) Grp 11, Pd 4      (C) Grp 17, Pd 5      (D) Grp 14, Pd 2

**UNIT 5 REVIEW WORKSHEET**

**Matching**

Alkali metals	Alkaline earth metals	Noble gases
Transition metals	Halogens	

- The \_\_\_\_\_ have a single electron in the highest energy level.
- The \_\_\_\_\_ achieve the electron configurations of noble gases by losing two electrons.
- The \_\_\_\_\_ achieve the electron configuration of noble gases by gaining one electron.
- The \_\_\_\_\_ have full s and p orbitals in the highest occupied energy levels.
- The \_\_\_\_\_ are stable and unreactive.
- The \_\_\_\_\_ are highly reactive nonmetals and readily form compounds with metals.
- The \_\_\_\_\_ are metals that are more reactive than the transition elements but less reactive than the alkali metals.

---

Atomic radius	Decrease	Noble gases
Electronegativity	Ionization energy	Increase
Metals	Noble gas configuration	Shielding effect
Nonmetals	Metalloid	

- \_\_\_\_\_ is the energy required to remove an electron from an atom.
- The attraction of an atom for an additional electron is called \_\_\_\_\_.
- When they have a(n) \_\_\_\_\_, ions have a stable, filled outer electron level.
- Along with the increased distance of the outer electrons from the nucleus, the \_\_\_\_\_ of the inner electrons causes ionization energy to decrease going down a column of the Periodic Table.
- A low ionization energy is characteristic of a(n) \_\_\_\_\_.
- Ionization energies tend to \_\_\_\_\_ across periods of the periodic table.
- An element with an extremely high ionization energy is classified as a(n) \_\_\_\_\_.

15. The distance from the nucleus to the highest occupied energy level is known as \_\_\_\_\_.
16. The \_\_\_\_\_ do not have measured electronegativities since they do not commonly form compounds.

=====

**Predict the oxidation number based on the electron configuration shown.**

- |                           |                           |
|---------------------------|---------------------------|
| 17. $1s^2 2s^2 2p^6 3s^2$ | 18. $1s^2 2s^2 2p^6 3s^1$ |
| 19. $1s^2 2s^2 2p^6$      | 20. $1s^2 2s^2 2p^5$      |
| 21. $1s^2 2s^2 2p^1$      |                           |

=====

**Choose the location of the element with the higher ionization energy.**

- |                        |    |                    |
|------------------------|----|--------------------|
| 22. Period 2, Group 14 | or | Period 3, Group 13 |
| 23. Period 4, Group 2  | or | Period 5, Group 2  |
| 24. Period 1, Group 18 | or | Period 2, Group 1  |
| 25. Period 3, Group 17 | or | Period 3, Group 18 |
| 26. Period 3, Group 17 | or | Period 2, Group 17 |
| 27. Period 3, Group 17 | or | Period 3, Group 16 |

=====

**Arrange the element locations in order of increasing electronegativity.**

- |                            |                        |                        |
|----------------------------|------------------------|------------------------|
| 28. (A) Period 4, Group 13 | (B) Period 3, Group 13 | (C) Period 5, Group 13 |
| 29. (A) Period 4, Group 2  | (B) Period 4, Group 16 | (C) Period 4, Group 15 |
| 30. (A) Period 2, Group 16 | (B) Period 2, Group 17 | (C) Period 3, Group 16 |
| 31. (A) Period 3, Group 15 | (B) Period 2, Group 16 | (C) Period 4, Group 14 |

=====

**Choose the location of the element with the larger atomic radius.**

- |                        |    |                    |
|------------------------|----|--------------------|
| 32. Period 2, Group 1  | or | Period 4, Group 1  |
| 33. Period 4, Group 2  | or | Period 4, Group 16 |
| 34. Period 4, Group 13 | or | Period 2, Group 13 |
| 35. Period 2, Group 16 | or | Period 2, Group 14 |
| 36. Period 3, Group 17 | or | Period 4, Group 17 |
| 37. Period 2, Group 2  | or | Period 6, Group 2  |
| 38. Period 3, Group 14 | or | Period 3, Group 16 |


=====

**Multiple Choice**

39. In any \_\_\_\_, the number of electrons between the nucleus and the outer energy level is the same.  
 (A) period    (B) group    (C) both    (D) neither
40. In a \_\_\_\_, electron affinity values decreases as atomic number increases.  
 (A) period    (B) group    (C) both    (D) neither



41. The halogens are considered a \_\_\_\_.  
 (A) period    (B) group    (C) both    (D) neither



***On the Periodic Table, show the location of the element with the...***

- 42. lowest ionization energy
- 43. most nonmetallic properties
- 44. smallest atomic radius
- 45. highest electronegativity
- 46. largest atomic radius
- 47. highest ionization energy
- 48. most metallic properties

***Answer the following questions.***

- 49. Explain the relationship between the relative size of an ion to its atom and the charge on the ion.
- 50. Explain why noble gases are inert and do not form ions.
- 51. Why do elements in the same family generally have similar properties?
- 52. If element X is a very reactive nonmetal, then the element with atomic number X + 1 should have what properties?

UNIT 5 REVIEW/SUMMARY WORKSHEET

1										18
	2				13	14	15	16	17	
		3	---	12						
		*								
		**								

*																				
**																				

- 1.) Color the "s" block area red.
- 2.) Color the "p" block area blue.
- 3.) Color the "d" block area green.
- 4.) Color the "f" block area orange.
- 5.) Draw an X in the boxes that represent the unreactive elements
- 6.) Draw a diagonal line (from upper left to lower right) in the area that represents the very reactive nonmetals.
- 7.) Draw a diagonal line (from upper right to lower left) in the area that represents the very reactive metals.
- 8.) Draw a purple capital letter "R" with a circle around it at the location that represents the element with the largest atomic radius.
- 9.) Draw a blue capital letter "I" with a diamond around it at the location that represents the element with the highest ionization energy.
- 10.) Draw a dark green capital letter "E" with a triangle around it at the location that represents the element with the highest electronegativity/electron affinity.
- 11.) Outline in black the boxes where metalloids with more nonmetallic properties are located.
- 12.) Outline in red the boxes where metalloids with more metallic properties are located.
- 13.) Draw a star in the location that represents the most metallic element (or most reactive metal).
- 14.) Draw a heart in the location that represents the most nonmetallic element (or most reactive nonmetal).